No 24

No 2
Doc 2
A ~~No 50~~ Sanson
No 39 South 6th St

Dissertation

on Pneumonia Typhoides

by

Refered March 30th 1827
W. R. H.

John R. Taylor

of

Virginia.

one of
am in
kel) t
on the
disease
through
death o
attention
eminen
from w
arrestin
discour
have a
happi
of the f
cure the
most a
I should

# Pneumonia Typhoides

In compliance with one of the established requisitions of this school. I am induced (however incompetent to the task I may feel) to make as the subject of a Thesis, some observations on the history and most prominent features of a disease; which from its wide and desolating ravages throughout every section of our country, and from the dark obscurity in which it is veiled; has elicited the attention and investigations of some of our most eminent physicians, in endeavouring to determine from whence its origin, and what the best method of arresting its career. Mortifying to their pride and discouraging to research and observation, their investigations have as yet been but sparingly rewarded. But happily for society and consoling to the feelings of the profession; if its pathology has been withheld, its cure the result of experience, has been crowned with the most ample and satisfactory success

I should be guilty of the base and unpardonable crime

of plag
thing o
It is n
attained
devised
in poi
checkin
We are
epidem
in the
state o
Canada
States.
it was
exciting
fatalit
was m
fatal-
shape,
perfec

of plagiarism, were I to claim the suggestion of any thing original in relation to this subject. Indeed it is not to be expected, and as such, I shall have perfectly attained my end, if by a compilation of facts derived from the most respectable authority, I succeed in pointing out the most approved method of checking its progress or effecting a cure

We are told by our Professor on the practice, that this epidemic first made its appearance in New Hampshire in the year eighteen hundred and six, from which state it spread throughout New England, entered the Canadies, and finally passed through the United States. In the year eighteen hundred and thirteen it was for the first time discovered in this City, not exciting much alarm, nor attended with much fatality; but upon its return the next winter, it was more envenomed, and its attack more frequently fatal — assuming for itself such a variety of shape, as to have procured the appellation of a perfect Proteus. It appears from one source (North)

and pe
that c
temper
disease
strong
males
more th
from o
le but
Withou
sympt
ling al
most fre
and hea
lind o
eyes vo
of the g
much
very mu
Lethargy

and perhaps a proriality of those of New England, that children, women, young people, and those of a sanguine temperament and lax fibre, are more liable to this disease. Whereas from another (Miller of Va) the strong and robust were peculiarly liable to it — males more frequently than females, and adults more than children — But that the diseases spring from one or the same common origin, there can be but little doubt

Without additional remarks, I shall proceed to the symptoms. They are described by Dr Chapman as being always attended with prostration and debility, most frequently commencing with alternations of chills and heat, the skin dry and pale, or mottled, the face livid or bronzed, alae nasi contracted, forehead pallished, eyes wild and glassy, the whole countenance expressive of the greatest anxiety and distress, pulse usually of not much volume, but hard like a small cord, soon sinking very much; the head much affected, wandering delirium, lethargy or stupor; the attack sometimes extremely violent

and re
in their
In man
extreme
loving
inion,
extreme
a parch
unexam
strictu
regular
violen
dejecte
mandel
agitat
beholde
desire
there
fever,
in 14°. o

and sudden, persons cloven down by it, whilst engaged in their common employments and died instantly. In many instances it comes on with pain in the extremities, at first very violent, afterwards fugitive, fixing in the head, and occasioning dimness of vision, perhaps blindness, delirium, coma, paralysis, extreme languor both of body and mind, and with a parched tongue. These symptoms if not checked, increase with rapidity - vertigo attended with painful stricture across the forehead and eyes ensue, unconquerable vigilance or somnolency, muttering or uttering of violent rhapsodies. If in his senses, is extremely dejected, sighs frequently, complains of uncontrolable wandering of his imagination, with the greatest anxiety agitation and distress, exciting in the mind of the beholder the highest sympathy, and the most ardent desire to relieve him of his wretchedness. Sometimes there is no local determination, neither chill nor fever, though the pulse be very frequent, at about a 140 or 150 in a minute, with some oppression at the

poracco
This w
symp
appear
commu
all the
and f
pant
a flur
gastr
full a
the fir
pleur
a day
frequ
of the
but n
great
if no
and

praecordia and a slight tension of the forehead. This state is highly deceitful and dangerous, the symptoms of debility on a sudden making their appearance and destroying the patient. But a more common form of the disease is that ushered in with all the signs of pneumonic inflammation, a chill and fever, pain in the side, difficult respiration, panting, coughing and spitting of bloody mucous, a flushed countenance, and suffused eye, much gastric distress, with vomiting bile. The pulse voluminous full and strong, though soft and easily compressible with the finger — resembling in all its symptoms bilious pleurisy. These symptoms however do not for more than a day or two, or perhaps for a few hours — Such cases are frequent. There is usually at first a slight affection of the throat, the patient not seeming very ill, but suddenly respiration is much impeded, and great prostration takes place. The throat and fauces if now examined, present the appearance of a dark and brownish or mahogony colour, altogether unlike that

of acte
met w
prostra
winks,
clumsi
colour
destrey
the bod
(spotte
more t
which
preser
and c
Serup
perhap
atten
Of th
that p
mista
after

* For
sym

of active inflammation — This form was not unfrequently met with in certain sections of our southern states — prostration takes place, the mind as also the body sinks, the patient is distracted, or in heavy stertorous slumbers. The tongue and fauces are covered with a dark coloured crust — countenance expressively haggard and distressed, vibices or petechiae are sometimes seen over the body, from whence the disease received its name — (Spotted fever) they however occur but very rarely not more than once in an hundred cases. There are also cases in which the throat is the chief seat of the disease, presenting the usual appearances of common cold, and are sometimes struck all at once, with the eruption of fatal debility and oppression. Such perhaps were the most frequent to the South and attended with the most serious consequences.x

Of the Diagnosis — It is said and perhaps with truth that physicians and others, are much more liable to mistake other diseases for an epidemic, especially after the occurrence of a few cases, than an epidemic

x For a very perfect and full delineation of the different symptoms, see Hall's history of Spotted fever

for oth
which
must
it cou
by the
the gr
exhibi
mista
for the
typhi
swoon
apopl
cases
several
met
which
showers
percei
threatin
from

for other diseases. But if we consider the close resemblance which this in its several stages bears to many others; we must be readily convinced of the liability of having it confounded not only by the theoretical, but often by the practical physician. It must be obvious from the great variety of symptoms, which this disease exhibits, that there will be extreme liability in mistaking it for several others, more particularly for the cynanche maligna, scarlatina, common typhus, rheumatism, hydrocephalus internus, worms choera morbus, hysteria, mania, pleuritis, apoplexy, paralysis, urticaria, colic: &c That sporadic cases should be particularly liable to be mistaken for several others, must be obvious to us all; for we occasionally meet with such in certain sections of the country, in which this epidemic has never prevailed. The consequence however of such mistakes, must sooner or later be perceived, if we compare the very different mode of treatment pointed out by experience in this disease, from that adopted in analagous cases. The best directions

of is n
mistak
physic
with th
hemac
should
make
much
the pa
more d
to be
obscure
disclose
who if
under
its co
no me
of a p
with
of the

it is said which can be given for avoiding such mistakes, is to remind the young and inexperienced physician, diligently to compare the history of this, with those to which it bears a resemblance. By this he may generally be enabled to distinguish it. But should such cases occur in which we are unable to make a proper distinction, we are advised to prescribe such remedies alone, as will not prove destructive to the patient, even should he be labouring under the more dangerous disease. It is nevertheless seriously to be regretted, that this disease clothed in as much obscurity as it often is, has before its real character been disclosed, precipitated beyond the reach of medicine, some who if their diseases had been timely managed and well understood, would have had an issue far less serious in its consequences— Of the Prognosis. It is a matter of no minor importance to the feelings and reputation of a physician, that he be able as far as is commensurate with human foresight, to form a correct judgement, of the probable course and termination of most

disea
cautio
fiad
natu
attac
sensi
when
great,
pulse
moist
the b
distru
hope
attent
attac
a rev
vital
tiume
tongu
reje

diseases. But with this under consideration great caution is necessary, particularly for those who have had but a small experience of its deceitful nature. But from the severity or mildness of the attack, we may form some idea, of its probable result. When the symptoms assume a mild form; when the prostration of strength is not uncommonly great, some degree of energy yet remaining in the pulse, the surface not cold, the tongue continuing moist; or if dry not much shriveled, or not assuming the bloodless appearance, if deglutition is not much obstructed, and the stomach retains medicine; we may hope and with confidence, by careful and unremitting attentions to carry the patient safely through the attack. And even though the symptoms may assume a severer form, the prostration of strength greater, the vital spark spark nearly extinguished, the pulse tremulous, the surface cold to the touch, the torpid tongue moved with difficulty; and the stomach rejecting its contents; even if these symptoms

are w
succe
to the
and s
throug
on me
and s
and s
confi
we are
stoma
admir
deglut
manne
ysteri
coma
of vol
on the
ouned
There

are within the control of medicine, and we succeed in time, in restoring warmth and moisture to the surface, increase the energy of the pulse, and vomiting is subdued, so as to enable us, through the stomach to invigorate the system; we may also have assurances of arresting its progress. And perhaps restore to society a valuable member, and receive the grateful smiles of joy and confidence from all surrounding friends. But we are authorized to say, if on the other hand, the stomach perseveringly rejects every medicine administered, a paralysis of the throat prevents deglutition, if we are unable by any means to restore warmth to the surface, and energy to the arterial system, if a torpid insensibility prevail, deep coma shuts up the senses, and destroys the power of voluntary motion, the respiration difficult, or the melancholy sound of apoplectic stupor, salutes our ears, our fears must be great our hopes but few. There are some other symptoms, which it is necessary

to obse
with d
they m
they l
is alm
happe
admir
oma,
no fut
appear
and c
come
can s
gener
and t
the sa
said
dang
depen
dang

to observe — petechiae and vibices are always attended with danger, the darker the colour, and the earlier they make their appearance, the greater danger do they indicate; a bronzed appearance of the forehead is almost invariably a fatal symptom. It sometimes happens, that unless the proper remedies are timely administered, dark coloured petechiae, complete coma, obstinate vomiting, a cold surface, little or no pulse, and a rigidity of the limbs make their appearance within a few hours after the attack. And if the fortunate period for the prevention of some of these symptoms, is suffered to pass, a cure can seldom be realized. It is laid down as a general rule, the shorter the cold or sinking stage is, and the sooner the reaction of the system is incited, the safer is the patient. Delirium or mania are also said to be much less dangerous than coma, the danger however of all or any of these states, depends much upon the stage of the fever, more dangerous in the cold stage than after reaction

is fou
and e
to chem
Moth
remar
as har
pronou
death
and o
indus
pour
little
insolv
prera
excep
This
appear
temp
cause
and

is formed. But however alarming the symptoms, and ineffectual our remedies, we are instructed to cherish and keep steadfastly in our minds this Motto - Dum spirat speratimus, and apply our remaining resources, however few, with as much assiduity as having as yet accomplished nothing. By thus persevering we may sometimes succeed in robbing death of some of the fairest objects of his selection, and of proving to the world that the rewards of industry and humanity are not always withheld from us - Causes - Of the causes of this disease little is as yet ascertained - It is as all other epidemics involved in much obscurity. It commences and prevails during cold weather, and is with a few exceptions dissipated by the warmth of Spring. This epidemic is asthenic in its nature, and appears to be dependent upon a low state of temperature. The remote and predisponent causes must be such as debilitate the system, and these are supposed to be influenced by

essar
d mat
munic
ning
dong
hon th
tility
n order
he earth
nimal
of them
whet
atmosph
unkno
exciting
an im
system
stimu
and a
of the

the seasons. The seasons in which the ulcerous sorethroat and malignant fevers prevailed, are said to have been unusually cold and wet. Such have they been also during the prevalence of this epidemic. Dampness and coldness are esteemed as having a direct influence upon the human system, producing relaxation and debility of its powers. Such when existing in more than an ordinary degree, render the vegetable productions of the earth, less nutritious, and less suitable for sustaining animal life. Whether these causes have been sufficient of themselves to predispose the system to this disease, or whether from some morbid constitution of the atmosphere, or more probably from some latent and unknown cause, is as yet not fully decided. The exciting are better understood, and are such as have an immediate or direct tendency to debilitate the system — as fatigue, fear, grief, loss of rest, abuse of stimuli, wetness, cold, and I may add abstinence, and any and every thing that depresses the system — of these cold is is the most frequent. That this

disease
that the
induced
expansi
its o
compel
superv
That th
opposed
have n
suffice
proper
mind,
drinks
the st
unleg
Of the
bark
acice
stage

disease neither originates in, nor is spread by contagion, is at this time, I believe pretty generally conceded. Indeed it cannot be otherwise, if we regard the wide expanse and universality of its prevalence, as well as its observance of the general laws of epidemics, compelling all other diseases to acknowledge its supremacy and to put on its attire. Treatment— That the tonic and stimulating method of cure, as opposed to the debilitating plan is the correct one, I have no doubt, for there is testimony abundantly sufficient to prove this fact. (North) Of the regimen proper in this fever, the means are rest of body and mind, avoiding fear, fresh air, agreable cordial drinks, liquid nutritious diet; and such as suits the stomach of the patient, best taken warm unless the heat is above the natural standard— Of the means to support the strength of the patient— bark and bitters, fermented liquors, wine, sulphuric acid &c In the comatose or sinking ~~or sinking~~ stage, we should be assiduous in external remedies,

and ad
tumul
excite
establis
useful
dark, a
Secume
system,
who se
to open
themsel
a thing
of breath
time c
cases se
he ha
other
patient
as the
necessa

and administer with a liberal hand the diffusible stimulants – wine – opium – ether &c until reaction is excited, sweat produced, and the pulse invigorated. – a blister applied to the nape of the neck is also useful. When petechiae and vibices are discovered, bark, wine and opium, are the appropriate remedies. Accumulated faeces acting as a cause of irritation to the system, it becomes necessary that the bowels be evacuated pro re nata, the means should be gentle, and so timed as not to operate during the sinking stage. Cases have presented themselves in which the symptoms were indicative of a turgescence of blood in the lungs – as great difficulty of breathing, and oppression at the breast, at the same time exhibiting a strangulated countenance. In such cases venesection may be proper, though North says he has never seen such a case. The wine and the other stimuli, never should be stopped, until the patient is restored to health, but the quantity diminished as the symptoms subside. There is perhaps no rule more necessary when the patient is low, than never to

suffer
ordial
ds to
author
I have
to prov
and M
vacuc
and bl
Woodw
bleedin
were a
rags. I
that
laugh
of the
in the
Dr Cha
have b
proper

suffer him to remain long without taking something cordial and nourishing.

As to the propriety of this treatment, I shall quote authority sufficient of itself to establish the point. I however feel almost as though, I were endeavouring to prove that which was already conceded. In Danielson's and Mann's essays, the stomach and bowels were vacuated, followed by a liberal use of wine, bark and blisters - this practice proved unsuccessful. Dr Woodward remarks having no guide but experience, bleeding, vomiting, purging, sweating and stimulating were all tried, sweating gave the most relief. Dr Beston says, I took small bleedings from the two first patients that I saw, the bad effects of which, though not fatal, taught me afterwards, extreme caution in the use of the lancet, or in making any other evacuation in the commencement of the disorder. Hale as also Dr Chapman observes that two modes of treating this disease have been proposed. By one set of practitioners, the most profuse use of direct stimulants, as wine, brandy,

as found
commo
middle
ravage
the up
rite
overs p
hely, h
Trunk,
is adra
diaphor
remedie
The carb
agnes
to give
very h
hot to
ured,
and
But

carbonate of ammonia, camphor &c have been recommended, whilst by another the treating has been decidedly prefered. By a comparison of the two modes of managing the disease, they have not the slightest doubt of the superiority of the latter. Of the means resorted to, to excite perspiration, nothing answered so well as the Dover's powders, once in three hours, strong wine whey given freely, hot fomentations applied to the extremities, to the trunk, and to the armpits. If called in after the disease is advanced, or in the latter stage, combine with diaphoretics the intrepid employment of that class of remedies denominated cordial stimulants or incitants. The carbonate of ammonia is here, said to be the ipse agmen or the unicum remedium, which should be given with a fearless hand, not less than v or x grs every hour. As cooperating means, strong wine whey, hot toddy, or madeira wine should be copiously used. At this crisis turpentine has been recommended and with the greatest probability of advantage. But should these means fail, and there be a rapid

sinking of the pulse; we must apply as a dernier resort, to the surface in various parts, a decoction of cantharides and spts turpentine to excite vesication, or the more irritant rubefacients, as a mixture of cayenne pepper with brandy.— This is the practice in the simple forms of the disease. But in the bilious, pneumonic, and anginose cases, it seems to be pretty generally conceded, that some difference of treatment is required. Emetics have been prescribed and with eminent advantage. As well for the purpose of evacuating the stomach, as for making salutary impressions upon the system generally. To be serviceable it is advised, that they be repeated in some instances several times, and those of the most active kind— emetic tartar. The James's has been recommended, and is said to display, when largely given, the combined operation of an emetic, purgative, and diaphoretic. These are to be succeeded by mercurial purgatives. After these measures have been premised, and we have so far accomplished

our end; we then subject the case to the stimulating diaphoretics, as in the simple states of the disease. If there be topical congestions or pains blisters become the appropriate remedy — as in anginose or pneumonic affections, delirium or coma, and such as are large enough to embrace the neck, or cover the head or chest, as the case may indicate. The most leading and conspicuous peculiarity of this disease, seems to depend on its assuming for the most part, the congestive as contradistinguished from the inflammatory condition — the veins are chiefly concerned in the first, the arteries in the latter. Bloodletting here has proved injurious, serving only to reduce the quantity in the circulating vessels, without unloading the engorged organs — By cups much good has been done, followed by the vesicating applications. Such also is pretty near the practice pursued by Drs Murphy and Miller of Va. The latter however speaking of this disease, says it presented two very different aspects — the inflammatory and typhoid, such certainly appears to have been

the ca
Treatm
ce de
Clusi
Blood
and el
ven g
a diff
implan
mar
bilico
From
together
also th
quite
diaph
Blood
vice
met
the re

the case, should we judge from the two plans of treatment pursued by himself. The one which we should all be compelled to consider, as a bilious inflammatory pleurisy, was treated with the most and repeated bloodletting, followed by emetics, purgatives, blisters, and diaphoretics — with this practice his patients were generally cured. But with the other type quite a different practice was adopted. From the inflammatory part which this disease wore, many were disposed to consider it a common bilious remittent of an inflammatory type. From many circumstances however attendant, together with the great prostration of strength, and also their want of success; induced him to adopt quite a different practice — the stimulating diaphoretics and with success. As it respects bloodletting in the typhoid of this disease, I will quote his words — "The three first cases I met with of the pneumonic form, I bled freely, the result of which was a speedy termination

death
ment. a
stumb
deple
ing, an
shun
strict
motor

What
hich I
art of
where
and m
only se
it wou
ency, a
and s
again

in death, in consequence of which I laid aside my lancet. Fifteen cases came under my notice in a particular part of the country, which were treated upon the depleting plan and all died — twenty in an other, and I have authority to say that at least one hundred and fifty, died under it, in other districts. The ill effects of this practice is further corroborated by Dr Williamson (Med. Repository)

If it could be otherwise than a repetition of what has been already detailed. I would cite a case which I had under my immediate care, during the latter part of the last winter. Suffice it to say that it was ushered in, with all the symptoms of bilious pleurisy, and was at first treated as such. Bloodletting was only resorted once, and much more sparingly than it would otherwise have been; but for my having seen, and also been cautioned, by an experienced and successful practitioner of the neighbourhood, against the indiscriminate and free use of the

lancet
swalle
had in
few
transf
made o
ness.
bleedin
neither
follow
Doses
tea, in
potass
most
remedi
and
for m
heart
almos
one

lancet in the Influenza, which was of very general prevalence at that time, and under which the pulse had invariably sunk. About forty eight hours after the commencement of the attack, it was translated from a most violent and excessive grade of high action, to one of the lowest I have ever seen. During which interval only one small bleeding was performed, from which the pulse was neither reduced, nor did the pain remit; — followed by a dose of calomel — oil — blisters and Dovers powders, assisted by the free use of the elm tea, impregnated with a portion of the nitrate of potash. After its assuming the typhoid type, the most liberal use of tonic, stimulating and diaphoretic remedies were had recourse to. And by persevering and unwearied attentions both day and night for nearly a week. I had the indescribable and heartfelt satisfaction, of wrestling from the almost embraces of that fell monster Death, one whose case had been considered, by all

who saw him, beyond the reach of medical assistance. I cannot however conclude the outline of this very interesting case (and which has been the ~~cause of~~ the selection of this subject) without lending my humble tribute of praise, to the wonderful efficacy, displayed by the combined operation of the Peruvian bark, the snake roots Polygala and Serpentaria — and the camphorated spirit